arXiv:1702.02625v2 [math.KT] 21 Feb 2017

ALGEBRAIC ATIYAH-SINGER INDEX THEOREM

NGUYEN LE DANG THI

Abstract. The aim of this work is to give an algebraic weak version of the Atiyah-Singer index theorem. We compute then a few small examples with the elliptic differential operator of order ≤ 1 coming from the Atiyah class in $\mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{O}_X, \Omega^1_{X/k})$, where $X \longrightarrow \mathrm{Spec}(k)$ is a smooth projective scheme over a perfect field k.

We follow Grothendieck [EGA4, §16.8] to recall briefly the notion of differential operators. Let $f : X \longrightarrow S$ be a morphism of schemes. Consider the Cartesian square



The diagonal $\Delta_{X/S} : X \longrightarrow X \times_S X$ is an immersion. One defines the n-th normal invariant of $\Delta_{X/S}$ as

$$\mathcal{P}^n_{X/S} = \Delta^{-1}_{X/S}(\mathcal{O}_{X\times_S X})/I_f^{n+1}.$$

It is clear that $\{\mathcal{P}^n_{X/S}\}_n$ form a projective system. One defines

$$\mathcal{P}^\infty_{X/S} = \varprojlim_n \mathcal{P}^n_{X/S}.$$

The first projection $p : X \times_S X \longrightarrow X$ induces an $\mathcal{O}_X$-algebra structure on $\mathcal{P}^n_{X/S}$ and the second projection $q : X \times_S X \to X$ induces a morphism

$$d^n_{X/S} : \mathcal{O}_X \longrightarrow \mathcal{P}^n_{X/S}.$$

If $\mathcal{F} \in \mathcal{O}_X - Mod$ is an $\mathcal{O}_X$-module, one defines

$$\mathcal{P}^n_{X/S}(\mathcal{F}) = \mathcal{P}^n_{X/S} \otimes_{\mathcal{O}_X} \mathcal{F}.$$

Let $\mathcal{F}, \mathcal{G} \in \mathcal{O}_X - Mod$ be two $\mathcal{O}_X$-modules. Let $D : \mathcal{F} \longrightarrow \mathcal{G}$ be a morphism of the underlying abelian sheaves. D is called a differential operator of order $\leq n$ relative S, if there exists a

Date: 06. 02. 2017.
1991 *Mathematics Subject Classification.* 14F22, 14F42.
Key words and phrases. K-theory, motivic cohomology, differential operators.

morphism of $\mathcal{O}_X$-modules $u : \mathcal{P}^n_{X/S}(\mathcal{F}) \to \mathcal{G}$, such that the following diagram commutes:



The morphism $D : \mathcal{F} \to \mathcal{G}$ between the underlying abelian sheaves is not $\mathcal{O}_X$-linear, but it is $f^{-1}\mathcal{O}_S$-linear. Let us denote by $\mathrm{Diff}^n_{X/S}(\mathcal{F}, \mathcal{G})$ the set of all differential operator of order $\leq n$ between $\mathcal{F}$ and $\mathcal{G}$. One defines

$$\mathrm{Diff}_{X/S}(\mathcal{F}, \mathcal{G}) = \bigcup_n \mathrm{Diff}^n_{X/S}(\mathcal{F}, \mathcal{G}).$$

It follows from the definition [EGA4, 16.8.3.1] that one has an isomorphism of abelian groups

$$\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{P}^n_{X/S}(\mathcal{F}), \mathcal{G}) \xrightarrow{\cong} \mathrm{Diff}^n_{X/S}(\mathcal{F}, \mathcal{G}).$$

If $f : X \to S$ is locally of finite type, then $\mathcal{P}^n_{X/S}$ is a quasi-coherent $\mathcal{O}_X$-module of finite type (cf. [EGA4, Prop. 16.3.9]). If $f : X \to S$ is locally of finite presentation, then $\mathcal{P}^n_{X/S}$ is a quasi-coherent $\mathcal{O}_X$-module of finite presentation (cf. [EGA4, Cor. 16.4.22]). Consequently, if $\mathcal{F}$ is a quasi-coherent $\mathcal{O}_X$-module of finite type resp. locally of finite presentation and $f : X \to S$ is locally of finite type resp. locally of finite presentation, then $\mathcal{P}^n_{X/S}(\mathcal{F}) = \mathcal{P}^n_{X/S} \otimes_{\mathcal{O}_X} \mathcal{F}$ is also a quasi-coherent $\mathcal{O}_X$-module of finite type resp. locally of finite presentation. If S is locally Noetherian and $f : X \to S$ is proper, then by [EGA3, Thm. 3.2.1] the higher direct image sheaf

$$R^q f_*(\mathcal{F}) = a_{Zar}(U \mapsto H^q_{Zar}(f^{-1}U, \mathcal{F}))$$

is coherent for any $\mathcal{F} \in Coh(X)$ and $\forall q \geq 0$. Consequently, if $X \longrightarrow \mathrm{Spec}(A)$ is proper over a Noetherian ring A, then for all coherent sheaf $\mathcal{F} \in Coh(X)$ the Zariski cohomology groups $H^q_{Zar}(X, \mathcal{F})$ are A-modules of finite type $\forall q \geq 0$. A differential operator $D : \mathcal{F} \longrightarrow \mathcal{G}$ induces a homomorphism of abelian groups

$$\Gamma_{ab}(X, \mathcal{F}) \overset{defn}{=} \mathrm{Hom}_{\underline{Ab}}(\mathbb{Z}, \mathcal{F}) \longrightarrow \mathrm{Hom}_{\underline{Ab}}(\mathbb{Z}, \mathcal{G}) \overset{defn}{=} \Gamma_{ab}(\mathbb{Z}, \mathcal{G}),$$

where we write $\underline{Ab}$ for the category of abelian sheaves on X and $\mathbb{Z}$ is the terminal object in $\underline{Ab}$. In general, we can not say anything about the kernel or cokernel of this homomorphism. However, D induces also a homomorphism of $\Gamma_{mod}(X, \mathcal{O}_X)$-modules

$$\Gamma_{mod}(X, \mathcal{P}^n_{X/S} \otimes_{\mathcal{O}_X} \mathcal{F}) = \mathrm{Hom}_{\mathcal{O}_X - Mod}(\mathcal{O}_X, \mathcal{P}^n_{X/S} \otimes_{\mathcal{O}_X} \mathcal{F}) \longrightarrow \mathrm{Hom}_{\mathcal{O}_X - Mod}(\mathcal{O}_X, \mathcal{G}) = \Gamma_{mod}(X, \mathcal{G}).$$

This motivates us to give the following definition:

Definition 1. (1) *Let $X \longrightarrow \mathrm{Spec}(k)$ be a k-scheme of finite type, where k is a field and $D : \mathcal{F} \to \mathcal{G}$ be a differential operator between two $\mathcal{O}_X$-modules of order $\leq n$. D is called a Fredholm operator, if the k-linear morphism between k-vector spaces*

$$D(X) : \Gamma_{mod}(X, \mathcal{P}^n_{X/k} \otimes_{\mathcal{O}_X} \mathcal{F}) \longrightarrow \Gamma_{mod}(X, \mathcal{G})$$

has finite-dimensional kernel and cokernel.

(2) *Let $X \to \mathrm{Spec}(A)$ be a scheme of finite type over a Noetherian ring A and $D : \mathcal{F} \to \mathcal{G}$ be a differential operator between two $\mathcal{O}_X$-modules. D is called a Fredholm operator, if the kernel and cokernel of the A-module morphism*

$$D(X) : \Gamma_{mod}(X, \mathcal{P}^n_{X/A} \otimes_{\mathcal{O}_X} \mathcal{F}) \longrightarrow \Gamma_{mod}(X, \mathcal{G})$$

are A-modules of finite type.

For a proper A-scheme X, where A is a Noetherian ring, any differential operator $D : \mathcal{F} \longrightarrow \mathcal{G}$ between coherent $\mathcal{O}_X$-modules is a Fredholm operator.

Definition 2. *Let X be a proper k-scheme and $D : \mathcal{F} \longrightarrow \mathcal{G}$ be a differential operator between coherent $\mathcal{O}_X$-modules. We define the index of D to be*

$$Ind(D) \stackrel{defn}{=} dim_k(Ker(D(X)) - dim_k(Coker(D(X)).$$

Let X be any scheme and $QsCoh(X)$ be the category of quasi-coherent $\mathcal{O}_X$-modules. Following Grothendieck [EGA2, §1.7] we define a vector bundle associated to a quasi-coherent sheaf $\mathcal{E} \in QsCoh(X)$ to be the X-scheme

$$\mathbb{V}(\mathcal{E}) \stackrel{defn}{=} \mathrm{Spec}(Sym_{\mathcal{O}_X}(\mathcal{E})).$$

The structure morphism $\pi : \mathbb{V}(\mathcal{E}) \to X$ is an affine morphism. For any morphism $f : Y \to X$, we have

$$\mathrm{Hom}_{Sch/X}(Y, \mathbb{V}(\mathcal{E})) \stackrel{\cong}{\longrightarrow} \mathrm{Hom}_{\mathcal{O}_X - Alg}(Sym_{\mathcal{O}_X}(\mathcal{E}), f_*\mathcal{O}_Y) \stackrel{\cong}{\longrightarrow} \mathrm{Hom}_{\mathcal{O}_X - Mod}(\mathcal{E}, f_*\mathcal{O}_Y).$$

We call $s_0 : X \to \mathbb{V}(\mathcal{E})$ the 0-section, if it corresponds to the 0-homomorphism $\mathcal{E} \to \mathcal{O}_X$. Now we assume $\mathcal{E}$ is an $\mathcal{O}_X$-module of finite type. Then $\pi : \mathbb{V}(\mathcal{E}) \to X$ is a morphism of finite type [EGA2, Prop. 1.7.11 (ii)]. So if X is a Noetherian scheme, then $\mathbb{V}(\mathcal{E})$ is also a Noetherian scheme. Consider the locally small abelian category $Coh(\mathbb{V}(\mathcal{E}))$ of coherent sheaves on $\mathbb{V}(\mathcal{E})$. We let

$$K_0(\mathbb{V}(\mathcal{E})) \stackrel{defn}{=} K(Coh(\mathbb{V}(\mathcal{E})))$$

to be the Grothendieck group of $\mathbb{V}(\mathcal{E})$. By [EGA2, Prop. 1.7.15] the morphism $s_0 : X \longrightarrow \mathbb{V}(\mathcal{E})$ is a closed immersion. Let $j : \mathbb{V}(\mathcal{E}) - s_0(X) \longrightarrow \mathbb{V}(\mathcal{E})$ be the complement open immersion. By [SGA6, §IX, Prop. 1.1] one has a localization exact sequence

$$K_0(X) \stackrel{s_{0*}}{\longrightarrow} K_0(\mathbb{V}(\mathcal{E})) \stackrel{j^*}{\longrightarrow} K_0(\mathbb{V}(\mathcal{E}) - s_0(X)) \longrightarrow 0. \tag{1}$$

If $f : X \to S$ is the structure morphism of X over a base scheme S, then any differential operator $D \in \mathrm{Diff}^n_{X/S}(\mathcal{F}, \mathcal{G})$ of order $\leq n$ between $\mathcal{O}_X$-modules defines a two terms complex of the underlying abelian sheaves

$$0 \longrightarrow \mathcal{F} \stackrel{D}{\longrightarrow} \mathcal{G} \longrightarrow 0.$$

Or equivalently, a two terms complex of $\mathcal{O}_X$-modules

$$0 \longrightarrow \mathcal{P}^n_{X/S} \otimes_{\mathcal{O}_X} \mathcal{F} \longrightarrow \mathcal{G} \longrightarrow 0.$$

We obtain then a two terms complex of $\mathcal{O}_{\mathbb{V}(\mathcal{E})}$-modules

$$0 \longrightarrow \pi^*(\mathcal{P}^n_{X/S} \otimes_{\mathcal{O}_X} \mathcal{F}) \longrightarrow \pi^*\mathcal{G} \longrightarrow 0.$$

Remark that if S is Noetherian and $f : X \longrightarrow S$ is of finite type, then X is also Noetherian, so $\pi^*(\mathcal{F})$, $\pi^*(\mathcal{P}^n_{X/S})$ and $\pi^*(\mathcal{G})$ are coherent $\mathcal{O}_{\mathbb{V}(\mathcal{E})}$-modules, if $\mathcal{F}$ and $\mathcal{G}$ are coherent. Moreover, by [EGA4, Prop. 16.4.5] there is a canonical isomorphism

$$\pi^*(\mathcal{P}^n_{X/S}) \xrightarrow{\cong} \mathcal{P}^n_{\mathbb{V}(\mathcal{E})/X}.$$

Definition 3. *Let $f : X \to S$ be a morphism of finite type, where S is a Noetherian scheme. Let $\mathcal{E} \in \mathcal{O}_X - Mod$ be an $\mathcal{O}_X$-module of finite type and $\pi : \mathbb{V}(\mathcal{E}) \to X$ be the associated vector bundle. Let $D \in \mathrm{Diff}^n_{X/S}(\mathcal{F}, \mathcal{G})$ be a differential operator between coherent $\mathcal{O}_X$-modules of order $\leq n$. Then D is called an elliptic operator with respect to $\mathcal{E}$, if there is an isomorphism*

$$\sigma(D) : \mathcal{P}^n_{\mathbb{V}(\mathcal{E})/X} \otimes_{\mathcal{O}_{\mathbb{V}(\mathcal{E})}} \pi^*\mathcal{F}_{|\mathbb{V}(\mathcal{E})-s_0(X)} \xrightarrow{\cong} \pi^*\mathcal{G}_{|\mathbb{V}(\mathcal{E})-s_0(X)},$$

The isomorphism $\sigma(D)$ is called the symbol of D.

By the localization exact sequence 1, an elliptic operator D of order $\leq n$ with respect to an $\mathcal{O}_X$-module of finite type $\mathcal{E}$ defines a class $[\mathcal{P}^n_{\mathbb{V}(\mathcal{E})/X}] \cdot [\pi^*\mathcal{F}] - [\pi^*\mathcal{G}] \in K_0(\mathbb{V}(\mathcal{E}))$, which becomes 0 after restricting to the complement of the 0-section. If $f : X \to S$ a morphism locally of finite type, then the sheaf of relative differentials $\Omega^1_{X/S} = I_f/I_f^2$ is an $\mathcal{O}_X$-module of finite type.

Definition 4. *Let $X \to S$ be a morphism of finite type over a Noetherian scheme S. A differential operator $D \in \mathrm{Diff}_{X/S}(\mathcal{F}, \mathcal{G})$ between coherent $\mathcal{O}_X$-modules is called elliptic, if it is elliptic with respect to $\Omega^1_{X/S}$.*

Let X be a Noetherian scheme. Denote by $Vect(X)$ the locally small exact category of locally free sheaves of finite rank on X. We let

$$K^0(X) \stackrel{defn}{=} K(Vect(X)).$$

For an arbitrary morphism of Noetherian schemes $f : Y \to X$ one has a functorial pullback homomorphism

$$f^* : K^0(X) \longrightarrow K^0(Y), \quad [\mathcal{E}] \mapsto [f^*\mathcal{E}].$$

For a proper morphism $f : Y \longrightarrow X$ between Noetherian schemes there is a pushforward homomorphism

$$f_* : K_0(Y) \longrightarrow K_0(X), \quad \mathcal{F} \mapsto \sum_i (-1)^i [R^i f_* \mathcal{F}],$$

which is well-defined as an exact sequence of coherent $\mathcal{O}_Y$-modules

$$0 \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}'' \longrightarrow 0$$

gives rise to an exact sequence of coherent $\mathcal{O}_X$-modules

$$\cdots \longrightarrow R^i f_*(\mathcal{F}') \longrightarrow R^i f_*(\mathcal{F}) \longrightarrow R^i f_*(\mathcal{F}'') \longrightarrow R^{i+1} f_*(\mathcal{F}') \longrightarrow \cdots$$

The functoriality of the pushforward follows easily from the Leray-Grothendieck spectral sequence

$$E_2^{p,q} = R^p g_*(R^q f_* \mathcal{F}) \Rightarrow R^{p+q}(g \circ f)_* \mathcal{F}.$$

If X is a regular Noetherian scheme, then the canonical homomorphism induced by the obvious exact embedding of categories

$$K^0(X) \xrightarrow{\delta} K_0(X)$$

is an isomorphism, since every coherent sheaf on a regular scheme has a finite locally free resolution.

Let now S be an arbitrary scheme. We denote by $SH(S)$ the stable motivic homotopy category together with the formalism of six functors $(f^*, f_*, f_!, f^!, \wedge, \underline{\mathrm{Hom}})$ as in [Ay08], [CD12] and [Hoy14, Appendix C]. Let $\mathbf{E}$ be a motivic ring spectrum parameterized by schemes. We will assume $\mathbf{E}$ is stable by base change, i.e. for any morphism of schemes $f : X \longrightarrow Y$ there is an isomorphism of spectra in $SH(X)$

$$f^*\mathbf{E}_Y \xrightarrow{\cong} \mathbf{E}_X.$$

Let S be a scheme and $\mathbf{E}_S \in SH(S)$ be a motivic ring spectrum. The unit $\varphi_S : \mathbb{S}^0 \to \mathbf{E}_S$ gives rise to a class

$$[\varphi_S] \in \widetilde{\mathbf{E}}^{2,1}(\mathbb{P}^1_S) \cong \mathbf{E}^{0,0}(\mathbb{P}^1_S).$$

We have a tower of S-schemes given by the obvious embeddings

$$\mathbb{P}^1_S \longrightarrow \mathbb{P}^2_S \longrightarrow \cdots \longrightarrow \mathbb{P}^n_S \longrightarrow \cdots$$

Let $\mathbb{P}^\infty_S = colim_n \mathbb{P}^n_S$, which is an object in the pointed unstable motivic homotopy category of Morel-Voevodsky $Ho_{\mathbb{A}^1,\bullet}(S)$ [MV99]. Let $i : \mathbb{P}^1_S \longrightarrow \mathbb{P}^\infty_S$ be the obvious map in $Ho_{\mathbb{A}^1,\bullet}(S)$, which gives rise to a map $\Sigma^\infty(i) : \Sigma^\infty(\mathbb{P}^1_S)_+ \longrightarrow \Sigma^\infty(\mathbb{P}^\infty_S)_+$. $\mathbf{E}_S$ is called oriented, if there is a class

$$\widetilde{\mathbf{E}}^{2,1}(\mathbb{P}^\infty_S) \ni c_S : \Sigma^\infty(\mathbb{P}^\infty_S)_+) \longrightarrow \mathbf{E}_S \wedge S^{2,1},$$

such that $i^*(c_S) = [\varphi_S]$. We say $\mathbf{E}$ is oriented, if $\mathbf{E}_S$ is oriented for any scheme S and for any morphism $f : T \to S$ one has $f^*(c_S) = c_T$. If S is regular, by [MV99, §4, 1.15, 3.7] there is a canonical isomorphism $B\mathbb{G}_m \cong \mathbb{P}^\infty_S$ in $Ho_{\mathbb{A}^1,\bullet}(S)$, which gives to a canonical isomorphism

$$Pic(S) \xrightarrow{\cong} [S_+, B\mathbb{G}_m] \xrightarrow{\cong} [S_+, \mathbb{P}^\infty_S].$$

For an oriented motivic ring spectrum $\mathbf{E}$, one can define the first Chern class as

$$c_1 : Pic(S) \xrightarrow{\cong} [S_+, \mathbb{P}^\infty_S] \xrightarrow{\Sigma^\infty} \mathrm{Hom}_{SH(S)}(\Sigma^\infty S_+, \Sigma^\infty \mathbb{P}^\infty_S) \xrightarrow{(c_S)_*} \mathrm{Hom}_{SH(S)}(\Sigma^\infty S_+, \mathbf{E}_S \wedge S^{2,1}) \xrightarrow{=} \mathbf{E}^{2,1}(S).$$

Assume S is a regular scheme. Let $X \longrightarrow S$ be a smooth S-scheme. For a vector bundle $\mathbb{V}(\mathcal{E})$ associated to a locally free $\mathcal{O}_X$-module $\mathcal{E}$ of finite rank r, there is an isomorphism (cf. [NSO09, Thm. 2.11])

$$\bigoplus_{i=0}^{r-1} \mathbf{E}^{*-2i,*-i}(X) \xrightarrow{\cong} \mathbf{E}^{*,*}(\mathbb{P}(\mathcal{E})), \quad (x_0, \cdots, x_{r-1}) \mapsto \sum_{i=0}^{r-1} p^*(x_i) \cup c_1^i(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(-1)),$$

where $\mathbb{P}(\mathcal{E}) = \mathrm{Proj}(Sym_{\mathcal{O}_X}(\mathcal{E})) \xrightarrow{p} X$ is the projective bundle associated to $\mathcal{E}$ [EGA2, §4]. So $\mathbf{E}^{*,*}(\mathbb{P}(\mathcal{E}))$ is a free module over $\mathbf{E}^{*,*}(X)$ with the basis

$$\{1, c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(-1)), \cdots, c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(-1))^{r-1})\}.$$

So for a vector bundle $\mathbb{V}(\mathcal{E})$ of rank r one can define the higher Chern classes

$$\mathbf{E}^{2i,i}(X) \ni c_i(\mathbb{V}(\mathcal{E})), \quad \sum_{i=0}^{r} p^*(c_i(\mathbb{V}(\mathcal{E})) \cup (-c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(-1))^{r-i} = 0,$$

where one puts $c_0(\mathbb{V}(\mathcal{E})) = 1$ and $c_i(\mathbb{V}(\mathcal{E})) = 0$ for $i \notin [0, r]$. Recall that one has a homotopy cofiber sequence [MV99, §3]

$$\mathbb{P}(\mathcal{E}) \xrightarrow{i} \mathbb{P}(\mathcal{E} \oplus \mathcal{O}_X) \xrightarrow{q} Th_X(\mathbb{V}(\mathcal{E})),$$

where $Th_X(\mathbb{V}(\mathcal{E}))$ denotes the Thom space of the vector bundle $\mathbb{V}(\mathcal{E})$. The homotopy cofiber sequence induces a long exact sequence

$$\cdots \longrightarrow \mathbf{E}^{*,*}(Th_X(\mathbb{V}(\mathcal{E}))) \xrightarrow{q^*} \mathbf{E}^{*,*}(\mathbb{P}(\mathcal{E} \oplus \mathcal{O}_X)) \xrightarrow{i^*} \mathbf{E}^{*,*}(\mathbb{P}(\mathcal{E})) \longrightarrow \cdots$$

The projective bundle formula tells us that i^* is a split epimorphism, so $\mathbf{E}^{*,*}(Th_X(\mathbb{V}(\mathcal{E})))$ is a free module of rank 1 over $\mathbf{E}^{*,*}(X)$, which is just $Ker(i^*)$. The Thom class is the unique class in $\mathbf{E}^{2r,r}(Th_X(\mathbb{V}(\mathcal{E})))$

$$t_{\mathbb{V}(\mathcal{E})} = (q^*)^{-1}(\sum_{i=0}^{r} p^*(c_i(\mathbb{V}(\mathcal{E})) \cup (-c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E}\oplus\mathcal{O}_X)}(-1))^{r-i}),$$

where $p : \mathbb{P}(\mathcal{E} \oplus \mathcal{O}_X) \to X$ is the structure morphism. Let $s : X \longrightarrow \mathbb{P}(\mathcal{E} \oplus \mathcal{O}_X)$ be its section. The Thom isomorphism is given by

$$th : \mathbf{E}^{*-2r,*-r}(X) \xrightarrow{\cong} \mathbf{E}^{*,*}(Th_X(\mathbb{V}(\mathcal{E}))), \quad x \mapsto x \cup t_{\mathbb{V}(\mathcal{E})}.$$

Its inverse is the composition

$$\begin{array}{ccccc} \mathbf{E}^{*,*}(Th_X(\mathbb{V}(\mathcal{E})) & \xrightarrowtail{q^*} & \mathbf{E}^{*,*}(\mathbb{P}(\mathcal{E} \oplus \mathcal{O}_X)) & \xrightarrow{s^*} & \mathbf{E}^{*-2r,*-r}(X) \\ & & \cong\uparrow & & \\ & & \bigoplus_{i=0}^{r} \mathbf{E}^{*-2i,*-i}(X) & & \end{array}$$

which sends

$$\sum_{i=0}^{r} p^*(x_i) \cup (-c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E}\oplus\mathcal{O}_X)}(-1))^{r-i}) \mapsto x_r.$$

If ξ is the universal quotient bundle on $\mathbb{P}(\mathcal{E} \oplus \mathcal{O}_X)$, i.e. there is an exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}(\mathcal{E}\oplus\mathcal{O}_X)}(-1) \longrightarrow p^*(\mathcal{E} \oplus \mathcal{O}_X) \longrightarrow \xi \longrightarrow 0.$$

Then by Whitney sum formula we have

$$q^*(t_{\mathbb{V}(\mathcal{E})}) = c_r(\xi).$$

Now let $KGL_S \in SH(S)$ be the motivic ring spectrum representing the algebraic K-theory constructed in [Rio10], [CD12, §13]. If S is regular and $X \in Sm/S$ is a smooth S-scheme, then one has a natural isomorphism

$$\mathrm{Hom}_{SH(S)}(\Sigma^\infty X_+, KGL_S) \xrightarrow{\cong} K_0(X). \tag{2}$$

KGL is an oriented motivic ring spectrum. Indeed, by Bott periodicity one has an isomorphism

$$KGL^{0,0}(S) \cong K_0(S) \cong \widetilde{KGL}^{0,0}(\mathbb{P}^1_S, \infty).$$

The Bott element β is the image of 1 under this isomorphism and $\beta \in KGL^{-2,-1}(S)$. The orientation of KGL is given by

$$c^{KGL} \overset{defn}{=} \beta^{-1} \cdot (1 - [\mathcal{O}_{\mathbb{P}^\infty_S}(1)]) \in KGL^{2,1}(\mathbb{P}^\infty_S).$$

For any line bundle $L \in Pic(S)$, its first Chern class is

$$c_1^{KGL}(L) = \beta^{-1}(1 - L^\vee).$$

For an arbitrary morphism $f : T \to S$ of regular schemes one has a commutative diagram [CD12, §13.1]

$$\begin{array}{ccccc} \mathrm{Hom}_{SH(S)}(\mathbf{1}_S, KGL_S) & \longrightarrow & \mathrm{Hom}_{SH(T)}(f^*\mathbf{1}_S, f^*KGL_S) & = & \mathrm{Hom}_{SH(T)}(\mathbf{1}_T, KGL_T) \\ \cong\downarrow & & & & \downarrow\cong \\ K_0(S) & & \xrightarrow[f^*]{} & & K_0(T) \end{array}$$

Now assume $\mathcal{E}$ is a locally free sheaf of finite rank on a smooth scheme X over a regular base S. We have a homotopy cofiber sequence (cf. [MV99, §3])

$$\mathbb{V}(\mathcal{E}) - s_0(X) \longrightarrow \mathbb{V}(\mathcal{E}) \longrightarrow Th_X(\mathbb{V}(\mathcal{E})),$$

where $Th_X(\mathbb{V}(\mathcal{E}))$ denotes the Thom space of the vector bundle $\mathbb{V}(\mathcal{E})$. This homotopy cofiber sequence gives rise to an exact sequence

$$\begin{aligned} \cdots \longrightarrow \mathrm{Hom}_{SH(S)}(\Sigma^\infty Th_X(\mathbb{V}(\mathcal{E})), KGL_S) &\longrightarrow \mathrm{Hom}_{SH(S)}(\Sigma^\infty \mathbb{V}(\mathcal{E})_+, KGL_S) \\ &\longrightarrow \mathrm{Hom}_{SH(S)}(\Sigma^\infty(\mathbb{V}(\mathcal{E}) - s_0(X))_+, KGL_S) \end{aligned}$$

The natural isomorphism 2 gives rise to a commutative diagram with exact rows (cf. [CD12, §13.4, (K6a)]), where we abbreviate $[-,-]$ for $\mathrm{Hom}_{SH(S)}(-,-)$:

$$\begin{array}{ccccc} [Th_X(\mathbb{V}(\mathcal{E})), KGL_S] & \longrightarrow & [\mathbb{V}(\mathcal{E}), KGL_S] & \longrightarrow & [\mathbb{V}(\mathcal{E}) - s_0(X), KGL_S] \\ & & \cong\downarrow & & \downarrow\cong \\ K_0(X) & \longrightarrow & K_0(\mathbb{V}(\mathcal{E})) & \longrightarrow & K_0(\mathbb{V}(\mathcal{E}) - s_0(X)) \end{array}$$

The bottom row is the exact sequence 1. For a Noetherian scheme X and an $\mathcal{O}_X$-module of finite type $\mathcal{E}$ the functor

$$\pi_* : Coh(\mathbb{V}(\mathcal{E})) \longrightarrow Coh(X)$$

is exact, since π is an affine morphism, which implies that $R^i\pi_*\mathcal{F} = 0, \forall i > 0, \forall \mathcal{F} \in Coh(\mathbb{V}(\mathcal{E}))$. So we still can define the pushforward

$$\pi_* : K_0(\mathbb{V}(\mathcal{E})) \longrightarrow K_0(X).$$

Hence, one may try to define the topological index by applying the homomorphism

$$K_0(Th_X(\mathbb{V}(\Omega^1_{X/k}))) \longrightarrow K_0(\mathbb{V}(\Omega^1_{X/k}))$$

then composing with $f_*\pi_*$. However, $\pi : \mathbb{V}(\mathcal{E}) \to X$ is not necessary projective, so we may run into difficulties, when we apply later the Grothendieck-Riemann-Roch theorem. Now we restrict ourselves to the situation where $f : X \to \mathrm{Spec}(k)$ is a smooth proper scheme over a field k. $\Omega^1_{X/k}$ is a locally free $\mathcal{O}_X$-module of finite rank. Let $D : \mathcal{F} \to \mathcal{G}$ be an elliptic operator of order $\leq n$. Let $\pi : \mathbb{V}(\Omega^1_{X/k}) \longrightarrow X$ denotes the vector bundles associated to $\Omega^1_{X/k}$. The symbol $\sigma(D)$ defines then an element

$$[\mathcal{P}^n_{\mathbb{V}(\Omega^1_{X/k})/X}] \cdot [\pi^*\mathcal{F}] - [\pi^*\mathcal{G}] \in K_0(\mathbb{V}(\Omega^1_{X/k})),$$

which lies in the image of the homomorphism

$$K_0(Th_X(\mathbb{V}(\Omega^1_{X/k}))) \longrightarrow K_0(\mathbb{V}(\Omega^1_{X/k})).$$

Let us denote by $[\sigma(D)] \in K_0(Th_X(\mathbb{V}(\Omega^1_{X/k})))$ a class, which is mapped to

$$[\mathcal{P}^n_{\mathbb{V}(\Omega^1_{X/k})/X}] \cdot [\pi^*\mathcal{F}] - [\pi^*\mathcal{G}],$$

such that its image in $K_0(\mathbb{V}(\Omega^1_{X/k}) - s_0(X))$ is trivial. We call this class $[\sigma(D)]$ a symbol class associated to the symbol $\sigma(D)$.

Definition 5. *Let $f : X \longrightarrow \mathrm{Spec}(k)$ be a smooth proper k-scheme. Let $D : \mathcal{F} \longrightarrow \mathcal{G}$ be an elliptic operator, where $\mathcal{F}, \mathcal{G} \in Coh(X)$. The topological index of D is defined as*

$$ind_{top}(D) = f_*(th^{-1}([\sigma(D)])),$$

where

$$th^{-1} : K_0(Th_X(\mathbb{V}(\Omega^1_{X/k}))) \xrightarrow{\cong} K_0(X)$$

is the inverse Thom isomorphism of algebraic K-theory and

$$f_* : K_0(X) \to K_0(\mathrm{Spec}(k)) = \mathbb{Z}, \quad \mathcal{E} \mapsto \sum_i (-1)^i [H^i_{Zar}(X, \mathcal{E})] = \chi(\mathcal{E})$$

If S is any scheme, let $H\mathbb{Q}$ denote the Beilinson rational motivic cohomology ring spectrum in $SH(S)$ constructed in [Rio10], [CD12, Defn. 14.1.2]. Let

$$ch_t : KGL_{\mathbb{Q}} \xrightarrow{\cong} \vee_{i\in\mathbb{Z}} H\mathbb{Q} \wedge S^{2i,i}$$

be the total Chern character, which is an isomorphism of rational spectra, if $S = \mathrm{Spec}(k)$, where k is a perfect field [Rio10, Defn. 6.2.3.9, Rem. 6.2.3.10]. One has the following result due to Riou:

Theorem 6. [Rio10, Thm. 6.3.1]*(Grothendieck-Riemann-Roch) Let k be a perfect field. Let $f : X \to S$ be a smooth projective morphism of smooth k-schemes. There is a commutative diagram in $SH(S)$*

$$\begin{array}{ccc} \mathbf{R}f_\star KGL_{\mathbb{Q}} & \xrightarrow{\mathbf{R}f_\star(ch\cdot Td(T_f))} & \bigvee_{i\in\mathbb{Z}} \mathbf{R}f_\star H\mathbb{Q} \wedge S^{2i,i} \\ \downarrow{\scriptstyle f_\star} & & \downarrow{\scriptstyle f_\star} \\ KGL_{\mathbb{Q}} & \xrightarrow[ch_t]{} & \bigvee_{i\in\mathbb{Z}} H\mathbb{Q} \wedge S^{2i,i} \end{array}$$

As explained in [CD12, §13.7] $f_\star$ induces the usual pushforward f_* on K-theory. Now we can compute

Proposition 7. *Let $f : X \to \mathrm{Spec}(k)$ be a smooth projective scheme over a perfect field k. Let $D : \mathcal{F} \to \mathcal{G}$ be an elliptic operator between coherent $\mathcal{O}_X$-modules. Then one has a formula*

$$ind_{top}(D) = \int_X ch(th^{-1}([\sigma(D)])) \cup Td(T_{X/k}),$$

where $\int_X$ means that we take the pushforward on motivic cohomology

$$\int_X : H^{2*,*}_{\mathcal{M}}(X, \mathbb{Q}) \longrightarrow H^{2*-2dim(X),*-dim(X)}_{\mathcal{M}}(\mathrm{Spec}(k), \mathbb{Q}) = \mathbb{Q}.$$

Proof. This is a trivial consequence of Thm. 6, where $S = \mathrm{Spec}(k)$. □

Now let $D \in \mathrm{Diff}^n_{X/A}(\mathcal{F}, \mathcal{G})$ be an arbitrary differential operator of order $\leq n$ on a projective scheme $X \to \mathrm{Spec}(A)$ over a Noetherian ring A and $\mathcal{F}, \mathcal{G} \in Coh(X)$. As D gives rise to an $\mathcal{O}_X$-module homomorphism

$$\mathcal{P}^n_{X/A} \otimes_{\mathcal{O}_X} \mathcal{F} \longrightarrow \mathcal{G},$$

we may take the twisting

$$\mathcal{P}^n_{X/A} \otimes_{\mathcal{O}_X} \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_X(m) \longrightarrow \mathcal{G} \otimes_{\mathcal{O}_X} \mathcal{O}_X(m),$$

which in turn gives us a differential operator of order $\leq n$

$$D(m) : \mathcal{F}(m) \longrightarrow \mathcal{G}(m).$$

We call $D(m)$ a twisting of D.

Proposition 8. *Let $X \longrightarrow \mathrm{Spec}(k)$ be a smooth projective scheme over a perfect field k and $D : \mathcal{F} \longrightarrow \mathcal{G}$ be a differential operator of order $\leq n$ between coherent $\mathcal{O}_X$-modules. Then there exists a number N_0, such that for all $N \geq N_0$ one has*

$$Ind(D(N)) = \int_X ch(([\mathcal{P}^n_{X/k}] \cdot [\mathcal{F}] - [\mathcal{G}])(N)) \cup Td(T_{X/k}).$$

Proof. This is quite trivial. By the result of Serre (see e.g [EGA3, Thm. 2.2.1, Prop. 2.2.2]), there is a number n_0, such that for all $q > 0$ and all $a \geq n_0$

$$H^q_{Zar}(X, \mathcal{P}^n_{X/k} \otimes_{\mathcal{O}_X} \mathcal{F}(a)) = 0,$$

and a number m_0, such that for all $q > 0$ and all $b \geq m_0$

$$H^q_{Zar}(X, \mathcal{G}(b)) = 0.$$

We take $N_0 = max(n_0, m_0)$ to be the maximal of n_0 and m_0. We have trivially that for all $N \geq N_0$

$$Ind(D(N)) = dim_k \Gamma_{mod}(X, \mathcal{P}^n_{X/k} \otimes_{\mathcal{O}_X} \mathcal{F}(N)) - dim_k \Gamma_{mod}(X, \mathcal{G}(N)).$$

The proposition follows now easily from the Hirzebruch-Riemann-Roch theorem

$$dim_k \Gamma_{mod}(X, E(N)) = \int_X ch(E(N)) \cup Td(T_{X/k}).$$

□

Now we are ready to state the following algebraic weak form of the Atiyah-Singer index theorem

Theorem 9. *Let $X \longrightarrow \mathrm{Spec}(k)$ be a smooth projective scheme over a perfect field k and $D \in \mathrm{Diff}^n_{X/k}(\mathcal{F}, \mathcal{G})$ be an elliptic differential operator of order $\leq n$, where $\mathcal{F}, \mathcal{G} \in Coh(X)$ are coherent $\mathcal{O}_X$-modules. There exists a number N_0, such that for all $N \geq N_0$ there is an equality*

$$Ind(D(N)) = ind_{top}(D(N)).$$

Proof. It remains to prove that

$$th^{-1}([\sigma(D(N))]) = ([\mathcal{P}^n_{X/k}] \cdot [\mathcal{F}] - [\mathcal{G}])(N),$$

which means that we have to show there is a commutative diagram

$$\begin{array}{ccc} K_0(Th_X(\mathbb{V}(\Omega^1_{X/k}))) & \longrightarrow & K_0(\mathbb{V}(\Omega^1_{X/k})) \\ & \searrow^{th^{-1}} & \downarrow{\scriptstyle s_0^*} \\ & & K_0(X) \end{array}$$

By construction, we have a commutative diagram

$$\begin{array}{ccc} K_0(Th_X(\mathbb{V}(\Omega^1_{X/k}))) & \longrightarrow & K_0(\mathbb{P}(\Omega^1_{X/k} \oplus \mathcal{O}_X)) \\ & \searrow^{th^{-1}} & \downarrow{\scriptstyle s^*} \\ & & K_0(X) \end{array}$$

So it remains to see that we have a commutative diagram

$$\begin{array}{ccc} K_0(\mathbb{V}(\Omega^1_{X/k})) & \longrightarrow & K_0(\mathbb{P}(\Omega^1_{X/k} \oplus \mathcal{O}_X)) \\ {\scriptstyle s_0^*}\downarrow & & \downarrow{\scriptstyle s^*} \\ K_0(X) & = & K_0(X) \end{array}$$

But this is quite obvious, since the closed immersion $s : X \longrightarrow \mathbb{P}(\Omega^1_{X/k} \oplus \mathcal{O}_X)$ is the composition of the 0-section $s_0 : X \longrightarrow \mathbb{V}(\Omega^1_{X/k})$ and $\mathbb{V}(\Omega^1_{X/k}) \longrightarrow \mathbb{P}(\Omega^1_{X/k} \oplus \mathcal{O}_X)$ (cf. [EGA2, Prop. 8.3.2]). □

Finally, we will give in the last part of this paper a few small examples. For this purpose, we always restrict now ourselves to the case of a smooth projective k-scheme $X \longrightarrow \mathrm{Spec}(k)$ of dimension $\dim(X) = d$, where k is a perfect field. Let us begin with the following lemma:

Lemma 10. *(de Jong and Starr* [dJS06, Lem. 2.1]*) Let $X \subset \mathbb{P}^n$ be a smooth complete intersection of type $(d_1, \cdots, d_c)$. Then*

$$ch(T_X) = n - c + \sum_{i=1}^{n-c}(n + 1 - \sum_{i=1}^{c} d_i^k)\frac{c_1(\mathcal{O}_X(1))^k}{k!}.$$

Proof. The proof is so elementary, so we reproduce the proof. Let $i : X \longrightarrow \mathbb{P}^n$ denote the closed embedding. Consider the short exact sequence

$$0 \longrightarrow T_X \longrightarrow i^*T_{\mathbb{P}^n} \longrightarrow \bigoplus_{i=1}^{c} \mathcal{O}_X(d_i) \longrightarrow 0.$$

Therefore,

$$ch(T_X) = ch(i^*T_{\mathbb{P}^n}) - c - \sum_{k=1}^{n-c}\sum_{i=1}^{c} d_i^k \frac{c_1(\mathcal{O}_X(1))^k}{k!}.$$

Consider the Euler sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^n} \longrightarrow \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus(n+1)} \longrightarrow T_{\mathbb{P}^n} \longrightarrow 0.$$

One has

$$ch(T_{\mathbb{P}^n}) = n + \sum_{k=1}^{n}(n+1)\frac{c_1(\mathcal{O}_{\mathbb{P}^n}(1))^k}{k!}.$$

The lemma follows easily. □

Consider the Atiyah class in $\mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{O}_X, \Omega^1_{X/k})$

$$0 \longrightarrow \Omega^1_{X/k} \longrightarrow \mathcal{P}^1_{X/k} \xrightarrow{u} \mathcal{O}_X \longrightarrow 0.$$

The morphism $u : \mathcal{P}^1_{X/k} \longrightarrow \mathcal{O}_X$ defines a differential operator of order ≤ 1

$$D_u \in \mathrm{Diff}^1_{X/k}(\mathcal{O}_X, \mathcal{O}_X).$$

Lemma 11. *D_u is an elliptic operator.*

Proof. Let $\pi : \mathbb{V}(\Omega^1_{X/k}) \longrightarrow X$ be the vector bundle associated to $\Omega^1_{X/k}$. We pullback the Atiyah class via π to obtain an exact sequence

$$0 \longrightarrow \Omega^1_{\mathbb{V}(\Omega^1_{X/k})/X} \longrightarrow \mathcal{P}^1_{\mathbb{V}(\Omega^1_{X/k})/X} \longrightarrow \mathcal{O}_{\mathbb{V}(\Omega^1_{X/k})} \longrightarrow 0.$$

For each point $P \in \mathbb{V}(\Omega^1_{X/k})$ we have by [EGA4, Cor. 16.4.12]

$$\mathcal{P}^1_{\mathbb{V}(\Omega^1_{X/k})/X,P} = \mathcal{O}_{\mathbb{V}(\Omega^1_{X/k}),P}/\mathfrak{m}^2_P.$$

The morphism $\pi^*(u) : \mathcal{P}^1_{\mathbb{V}(\Omega^1_{X/k})/X} \longrightarrow \mathcal{O}_{\mathbb{V}(\Omega^1_{X/k})}$ becomes obviously an isomorphism at each point $P \in \mathbb{V}(\Omega^1_{X/k}) - s_0(X)$. □

From now on we always consider the elliptic differential operator D_u. We write $H = c_1(\mathcal{O}_X(1))$ and for a curve X

$$deg(H) = \int_X c_1(\mathcal{O}_X(1)).$$

Proposition 12. *Let $C \subset \mathbb{P}^3$ be a smooth curve of genus 1, which is a smooth complete intersection of two quadrics. For all integer number $n \geq 0$ one has*

$$ch(\Omega^1_{C/k}(n)) \cup Td(T_{C/k}) \in H^{2*,*}_{\mathcal{M}}(C, \mathbb{Z}).$$

For $N >> 0$ one has

$$Ind(D_u(N)) = 4N.$$

Proof. C is smooth complete intersection of two quadrics. By the lemma 10 we have $ch(\Omega^1_{C/k}) = 1$. This implies

$$ch(\Omega^1_{C/k}(n)) = ch(\mathcal{O}_C(n)) = (1 + n \cdot H),$$

for any number $n \geq 0$, where we write $H = c_1(\mathcal{O}_C(1))$ for the hyperplane section. As $g_C = 1$, the canonical divisor $K_C = 0$ is trivial. So we have $Td(T_{C/k}) = 1$ and hence

$$ch(\Omega^1_{C/k}(n)) \cup Td(T_{C/k}) = (1 + n \cdot H) \in H^{2*,*}_{\mathcal{M}}(C, \mathbb{Z}).$$

This implies easily that $Ind(D_u(N)) = N \cdot deg(H) = 4N$ for $N >> 0$. □

Proposition 13. *Let $X \longrightarrow \mathbb{P}^1$ be a flat projective morphism over a perfect field K, such that the generic fiber X_η is a smooth curve of genus one in $\mathbb{P}^2$. Then there exists a number $N >> 0$, such that*

$$Ind(D_u) \equiv 1 \mod 2.$$

Proof. We have for $N >> 0$:

$$Ind(D_u(N)) = dim_K \Gamma_{mod}(X, \Omega^1_{X/K}(N)).$$

We take two embeddings $i_1 : \mathbb{P}^1 \longrightarrow \mathbb{P}^M$ and $i_2 : X \longrightarrow \mathbb{P}^M$, such that they are compatible with f

$$\begin{array}{ccc} X & \xrightarrow{f} & \mathbb{P}^1 \\ & \searrow^{i_2} & \downarrow{\scriptstyle i_1} \\ & & \mathbb{P}^M \end{array}$$

Let $N_0 >> 0$ be a big integer number. By [EGA3, Cor. 7.9.13, Seconde partie] $f_*(\Omega^1_{X/K}(N_0))$ is a locally free $\mathcal{O}_{\mathbb{P}^1}$-module of rank $\chi(X_\eta, \Omega^1_{X_\eta,\eta}(N_0))$. By flat base change for Zariski cohomology of coherent sheaves it is enough to compute $\chi(X_{\bar{\eta}}, \Omega^1_{X_{\bar{\eta}}}(N_0))$, where $X_{\bar{\eta}}$ is the base change of X_η to an algebraic closure $\bar{\eta}$. $X_{\bar{\eta}}$ must be an elliptic curve. For an elliptic curve (E, P_0), we know that $\Omega^1_E \cong \mathcal{O}_E$ and $\mathcal{L}(3P_0)$ is very ample, i.e. $\mathcal{L}(3P_0) \cong \mathcal{O}_E(1)$. The last isomorphism means simply that we can embed

$$|3P_0| : E \longrightarrow \mathbb{P}^2.$$

By Riemann-Roch theorem for curves we also have $dim\, H^0(E, \mathcal{L}(nP_0)) = n$. Now we can apply the theorem of Grothendieck for the decomposition of vector bundles on $\mathbb{P}^1$ and we have

$$f_*(\Omega^1_{X/K}(N_0)) \cong \bigoplus_{i=1}^{3N_0} \mathcal{O}_{\mathbb{P}^1}(a_i), \quad a_i \in \mathbb{Z}.$$

Let $N_0' >> 0$ be another big integer number. We have

$$f_*(\Omega^1_{X/K}(N_0))(N_0') \cong \bigoplus_{i=1}^{3N_0} \mathcal{O}_{\mathbb{P}^1}(a_i + N_0').$$

So

$$dim_K f_*(\Omega^1_{X/K}(N_0))(N_0') = \sum_{i=1}^{3N_0}(1 + a_i + N_0') = 3N_0(N_0' + 1) + \sum_{i=0}^{3N_0} a_i.$$

If this dimension is already odd, then there is nothing to prove. So we assume

$$3N_0(N_0' + 1) + \sum_{i=1}^{3N_0} a_i \equiv 0 \mod 2.$$

Let $N_0'' >> 0$ be another big integer number. After twisting by $\mathcal{O}(N_0'')$ we have

$$dim_K f_*(\Omega^1_{X/K}(N_0))(N_0' + N_0'') = 3N_0(N_0' + N_0'' + 1) + \sum_{i=0}^{3N_0} a_i.$$

Now we take simply $N_0 \equiv 1 \mod 2$, $N_0' \equiv 1 \mod 2$ and $N_0'' \equiv 0 \mod 2$. We obtain

$$dim_K f_*(\Omega^1_{X/K}(N_0))(N_0' + N_0'') \equiv 1 \mod 2$$

By projection formula we have for any number $n > 0$ an isomorphism

$$f_*(\Omega^1_{X/K}) \otimes_{\mathcal{O}_{\mathbb{P}^1}} \mathcal{O}_{\mathbb{P}^1}(n) \xrightarrow{\cong} f_*(\Omega^1_{X/K}) \otimes_{\mathcal{O}_{\mathbb{P}^1}} i_1^*\mathcal{O}_{\mathbb{P}^M}(n) \xrightarrow{\cong} f_*(\Omega^1_{X/K} \otimes_{\mathcal{O}_X} f^* i_1^* \mathcal{O}_{\mathbb{P}^M}(n))$$
$$\xrightarrow{\cong} f_*(\Omega^1_{X/K} \otimes_{\mathcal{O}_X} i_2^* \mathcal{O}_{\mathbb{P}^M}(n)) \xrightarrow{\cong} f_*(\Omega^1_{X/K} \otimes_{\mathcal{O}_X} \mathcal{O}_X(n)).$$

This implies

$$dim_K \Gamma_{mod}(X, \Omega^1_{X/K}(N_0 + N_0' + N_0'')) = dim_K \Gamma_{mod}(\mathbb{P}^1, f_*(\Omega^1_{X/K}(N_0 + N_0' + N_0''))) \equiv 1 \mod 2,$$

where the first equality follows easily from the adjunction

$$f^* : \mathcal{O}_{\mathbb{P}^1} - Mod \leftrightarrows \mathcal{O}_X - Mod : f_*.$$

Therefore, we can conclude that there exists a number $N = N_0 + N_0' + N_0'' >> 0$ such that $Ind(D_u(N)) \equiv 1 \mod 2$, which finishes the proof. □

Proposition 14. *Let $X \subset \mathbb{P}^4$ be a smooth $K3$-surface, which is a smooth complete intersection, over a perfect field k.*

(1) *If $deg(X) = 4$, then for a big number $N >> 0$ one has*
$$Ind(D_u(N)) = 4N^2 - 36N + 1.$$
(2) *If $deg(X) = 6$, then for a big number $N >> 0$ one has*
$$Ind(D_u(N)) = 6N^2 - 60N - 20.$$

Proof. We write $c_1(\mathcal{O}_X(1)) = H$ and K for the canonical divisor. For an algebraic surface one has

$$Td(T_{X/k}) = 1 - \frac{1}{2}K + \frac{1}{12}(K^2 + c_2).$$

For a smooth projective $K3$-surface one has $K = 0$ and $K^2 = 0$. $\Omega^1_{X/k}$ is the dual bundle of $T_{X/k}$. So we have by lemma 10

$$ch(\Omega^1_{X/k}) = 2 - (5 + d_1 + d_2) \cdot H + (5 - d_1^2 - d_2^2) \cdot \frac{H^2}{2}.$$

For any number $n \geq 0$ one has

$$ch(\mathcal{O}_X(n)) = 1 + n \cdot H + \frac{1}{2}n^2 \cdot H^2.$$

If $deg(X) = 4$ then $(d_1, d_2) = (2, 2)$ and if $deg(X) = 6$ then $(d_1, d_2) = (2, 3)$. The result follows now easily, since $deg(X) = \int_X H^2$ and $\int_X c_2 = 24$. □

References

[Ay08] J. Ayoub, Les six opération de Grothendieck et le formalisme des cycles évanescents dans le monde motivique I and II. Astérisque **314**, **315**, (2008).

[CD12] C. D. Cisinski, F. Déglise, Triangulated category of mixed motives, arXiv:0912.2110v3 [math.AG], Preprint (2012).

[dJS06] A. J. de Jong, J. Starr, Low degree complete intersections are rationally simply connected, Preprint, 74 pages.

[EGA2] A. Grothendieck, J. Dieudonné, Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes, Publ. Math. IHÉS **8** (1961).

[EGA3] A. Grothendieck, J. Dieudonné, Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents, Publ. Math. IHÉS **11** (1961) and **17** (1963).

[EGA4] A. Grothendieck, J. Dieudonné, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV, Publ. Math. IHÉS **20**, **24**, **28**, **32** (1964-1967).

[Hoy14] M. Hoyois, A quadratic refinement of the Grothendieck-Lefschetz-Verdier trace formula, Algebr. Geom. Topol. 14, no. **6**, (2014).

[MV99] F. Morel, V. Voevodsky, $\mathbb{A}^1$-homotopy theory of schemes, IHÉS Publ. Math., (**90**), 45-143, (1999).

[NSO09] N. Naumann, M. Spitzweck, Paul Arne Østvær, Chern classes, K-theory and Landweber exactness over nonregular base schemes, in Motives and Algebraic Cycles: A Celebration in Honour of Spencer J. Bloch, Fields Inst. Comm., Vol. **56**, (2009).

[Rio10] J. Riou, Algebraic K-theory, $\mathbb{A}^1$-homotopy and Riemann-Roch theorems, J. of Topology **3**, (2010), pp. 229-264.

[SGA6] P. Berthelot, A. Grothendieck, L. Illusie, Théorie des intersections et théorème de Riemann-Roch, Lect. notes in Math. **225**, Springer-Verlag, (1971).

E-mail address: nguyen.le.dang.thi@gmail.com